A GUIDE TO MONITORING FOR SIGNS AND SYMPTOMS

LEARN WHY IT'S IMPORTANT AND HOW TO DO IT

WHAT ARE IMFINZI AND IMJUDO?

IMFINZI and IMJUDO are prescription medicines used to treat adults with **a type of lung cancer called NSCLC**. IMFINZI may be used in combination with IMJUDO and chemotherapy that contains platinum when your NSCLC has spread to other parts of your body (metastatic) **and** your tumor does not have an abnormal "EGFR" or "ALK" gene.

It is not known if IMFINZI and IMJUDO are safe and effective in children.

SELECT SAFETY INFORMATION

What is the most important information I should know about IMFINZI and IMJUDO?

IMFINZI and IMJUDO are medicines that may treat certain cancers by working with your immune system. IMFINZI and IMJUDO can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended.

For Important Safety Information, please see pages 4-8.

Please see additional Important Safety Information throughout and click here for Full Prescribing Information including Medication Guide for <u>IMFINZI</u> and <u>IMJUDO</u>.





UNDERSTANDING SIDE EFFECTS

IMFINZI and IMJUDO are medicines that may treat certain cancers by working with your immune system. If you're starting treatment with IMFINZI + IMJUDO with chemotherapy, it's important for you to understand that you may experience side effects. IMFINZI and IMJUDO may cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become severe or even life-threatening and can lead to death. You can have more than one of these problems at the same time. These problems may happen anytime during treatment or even after your treatment has ended. That's why tracking your side effects is such an important part of your treatment.

MONITORING AND REPORTING SIGNS OF SIDE EFFECTS

TRACK HOW YOU FEEL NORMALLY

There's a chance you may already be feeling side effects from your cancer, previous treatments, or even other medications. So it's important to write down how you feel before treatment to help you monitor any changes.

CATCH YOUR SIDE EFFECTS EARLY

Getting medical treatment right away may help keep any problems you may experience from becoming more serious. Your healthcare provider will check you for these side effects during your treatment with IMFINZI + IMJUDO with chemotherapy. Your healthcare provider may also need to delay or completely stop treatment with IMFINZI + IMJUDO with chemotherapy if you have severe side effects.

STAY CONNECTED WITH YOUR CARE TEAM

It's important to **let your care team know** of any changes in how you're feeling. Talk to them about how you're currently feeling, new signs of side effects, or any other change. Remember, a small side effect can lead to something more serious if not treated.

Turn the page for an overview of potential urgent side effects. Use the pages in this brochure to track the signs of side effects.

If you have any questions about your treatment, ask your doctor. The information in this brochure does not take the place of talking to your care team about your Stage 4 non-small cell lung cancer (NSCLC) treatment.



For your safety, remember to fill out the wallet card enclosed in this brochure. Please see page 9.





URGENT SIDE EFFECTS

Look out for the following side effects listed below. Getting medical treatment right away may help keep these problems from becoming more serious. **Call or see your healthcare provider right away if you develop any of the following symptoms or if these symptoms worsen.**



Do not attempt to treat any of the side effects without talking to your care team first.

Call or see your healthcare provider right away if you develop any new or worsening signs or symptoms including:

Lung problems



- Cough Shortness of breath
- Shortness of bre
- Chest pain

Intestinal problems

- Diarrhea (loose stools) or more frequent bowel movements than usual
- Stools that are black, tarry, sticky, or have blood or mucus
- Severe stomach-area (abdomen) pain or tenderness

Liver problems

4

- Severe nausea or vomiting
- Pain on the right side of your stomach area (abdomen)

Yellowing of your skin or the whites of your eyes

- Dark urine (tea colored)
- Bleeding or bruising more easily than normal

Hormone gland problems



away or unusual headachesEye sensitivity to light

Headaches that will not go

- Eye problems
- Rapid heartbeat
- Increased sweating
- Extreme tiredness
- Weight gain or weight loss
- Feeling more hungry or thirsty than usual

Kidney problems

- Decrease in your amount of urine
- Blood in your urine
- Swelling of your ankles
- Loss of appetite

Skin problems



- Rash
- Itching
- Skin blistering or peeling
- · Painful sores or ulcers in mouth or nose, throat, or genital area
- Fever or flu-like symptoms
- Swollen lymph nodes

Pancreas problems



- Pain in your upper stomach area (abdomen)
- Severe nausea or vomiting
- Loss of appetite

durvalumab

Injection for Intravenous Use 50 mg/mL



Urinating more often than usua

Changes in mood or behavior,

such as decreased sex drive.

irritability, or forgetfulness

Your voice gets deeper

Dizziness or fainting

Hair loss

Feeling cold

Constipation

URGENT SIDE EFFECTS (cont'd)

Problems can also happen in other organs and tissues. These are not all of the signs and symptoms of immune system problems that can happen with IMFINZI and IMJUDO. Call or see your healthcare provider right away for any new or worsening signs or symptoms.

Signs or symptoms may include:



- Chest painIrregular heartbeats
- Shortness of breath or swelling of ankles
- Confusion
- Sleepiness
- Memory problems
- Changes in mood or behavior

- Stiff neck, balance problems, tingling, numbness or weakness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eye sight
- Persistent or severe muscle pain
- or weakness, muscle cramps, joint pain, joint stiffness or swelling
- Low red blood cells and bruising

Infusion reactions that can sometimes be severe or life-threatening.

Signs and symptoms of infusion reactions may include:

- Chills or shaking
- Itching or rash
- Flushing

Shortness of breath
 or wheezing

- Dizziness
 Feel like passing out
- Fever
- Back or neck pain

Complications, including graft-versus-host disease (GVHD), in people who have received a bone marrow (stem cell) transplant that uses donor stem cells (allogeneic). These complications can be serious and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with IMFINZI. Your healthcare provider will monitor you for these complications

ADDITIONAL SAFETY INFORMATION

Getting medical treatment right away may help keep these problems from becoming more serious. Your healthcare provider will check you for these problems during your treatment with IMFINZI and IMJUDO. Your healthcare provider may treat you with corticosteroid or hormone replacement medicines Your healthcare provider may also need to delay or completely stop treatment with IMFINZI and IMJUDO if you have severe side effects

Before you receive IMFINZI and IMJUDO, tell your healthcare provider about all of your medical conditions, including if you:

- have immune system problems such as Crohn's disease, ulcerative colitis, or lupus
- have received an organ transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)
- have received radiation treatment to your chest area
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. IMFINZI and IMJUDO can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider will give you a pregnancy test before you start treatment with IMFINZI and IMJUDO.
- You should use an effective method of birth control during your treatment and for 3 months after the last dose of IMFINZI and IMJUDO. Talk to your healthcare provider about birth control methods that you can use during this time.
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with IMFINZI and IMJUDO.
- are breastfeeding or plan to breastfeed. It is not known if IMFINZI and IMJUDO pass into your breast milk. Do not breastfeed during treatment and for 3 months after the last dose of IMFINZI and IMJUDO.

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.



ADDITIONAL SAFETY INFORMATION (cont'd)

WHAT ARE THE POSSIBLE SIDE EFFECTS OF IMFINZI AND IMJUDO?

IMFINZI and IMJUDO can cause serious side effects (see pages 4-6):

The most common side effects of IMFINZI when used in combination with IMJUDO and platinum-containing chemotherapy in adults with metastatic NSCLC include:

- Nausea
- Feeling tired or weak
- Muscle or bone pain
- Decreased appetite
- Rash

8

• Diarrhea

Tell your healthcare provider if you have any side effect that bothers you or that does not go away. These are not all the possible side effects of IMFINZI and IMJUDO. Ask your healthcare provider or pharmacist for more information.

Call your doctor for medical advice about side effects. You are encouraged to report negative side effects of AstraZeneca prescription drugs by calling 1-800-236-9933. If you prefer to report these to the FDA, either visit www.FDA.gov/medwatch or call 1-800-FDA-1088.

Please see additional Important Safety Information throughout and click here for Full Prescribing Information including Medication Guide for <u>IMFINZI</u> and <u>IMJUDO</u>.

INFORMATION IN YOUR POCKET

THE WALLET CARD

(2)

3

If you ever experience an emergency and need to see a doctor who's not on your regular care team, keeping this card in your wallet can let them know that you're receiving an immunotherapy and help them get in touch with your doctor, if they need to.

I am receiving immunotherapy.				
Please contact my oncologist immediately before treatment.				
My Name:				
Immunotherapy:				
Oncologist:				
Contact:				
AstraZeneca 🕏 ©2017 AstraZeneca. All rights reserved. 3298301 2/17				

3 TIPS TO MAKE SURE YOU GET THE APPROPRIATE CARE

- Once you've started treatment: Fill out this card and carry it with you at all times.
- Even after you stop treatment: Keep this card with you for at least 3 months, because side effects can still occur during that time.
- Take a picture: If you have a phone with a camera, take a picture of this card just in case you leave it at home.





HOW TO TRACK SIGNS OF SIDE EFFECTS

SIDE EFFECTS TRACKER

Below is a helpful guide you can use to track signs of side effects during a single day.

	BOWEL MOVEMENTS	SKIN PROBLEMS	MAL PAIN
What's normal for me	I usually have <u>3</u> bowel movements a day and they are <u>hard</u>	My skin problems flare up: © NEVER OFTEN	On a scale of 1 to 10 (10 = severe), my pain is usually: <u>4</u> and lasts: <u>3-5</u> hours
Date, appointments, etc	Write down how many bowel movements you've had today and other details	Today, my skin problems flared up: OOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOOO	How painful? How long? Describe your pain today
January 12 infusion at Dr. White	2, very loose stools	NEVER OFTEN Skin rash and itchiness	- Morning stomach pain - 4 in intensity - Lasted 3 hrs
		NEVER OFTEN	
		NEVER OFTEN	




It's important to write down how you normally feel before your first infusion, so you can notice any changes that may occur during treatment.

NAME:

TIREDNESS		OTHER SIGNS OF SIDE EFFECTS	WHAT ACTIVITIES DID YOU DO?
l usually sleep: hours a day	l usually have: 3 meals a day	I usually experience:	
Write down how many hours you've slept today	Write down how many times you've eaten today and other details	Write down how many times you felt this or other signs of side effects today	
5	Ate 3 meals today.		went to the park today.

SIDE EFFECTS TRACKER



NAME:

12.			
J TIREDNESS	APPETITE	OTHER SIGNS OF SIDE EFFECTS	WHAT ACTIVITIES DID YOU DO?
I usually sleep:	I usually have:	I usually experience:	
hours a day			
Nrite down how many ours you've slept today	Write down how many times you've eaten today and other details	Write down how many times you felt this or other signs of side effects today	





SIDE EFFECTS TRACKER

	BOWEL MOVEMENTS	SKIN PROBLEMS	MAL PAIN
What's normal for me	I usually have bowel movements a day and they are 	My skin problems flare up: OOOOO NEVER OFTEN	On a scale of 1 to 10 (10 = severe), my pain is usually: and lasts: hours
Date, appointments, etc	Write down how many bowel movements you've had today and other details	Today, my skin problems flared up: OOOOOO NEVER OFTEN Note other details	How painful? How long? Describe your pain today
		NEVER OFTEN	
		O-O-O-O-O NEVER OFTEN	

NAME:

TIREDNESS	APPETITE	OTHER SIGNS OF SIDE EFFECTS	WHAT ACTIVITIES DID YOU DO?
l usually sleep:	I usually have:	I usually experience:	
hours a day			
Write down how many hours you've slept today	Write down how many times you've eaten today and other details	Write down how many times you felt this or other signs of side effects today	









IMFINZI and IMJUDO are registered trademarks of the AstraZeneca group of companies. ©2024 AstraZeneca. All rights reserved. US-84435 Last Updated 1/24